

# Ep-CAM / CD326 (Rat) Antibody

Mouse Monoclonal Antibody [Clone EpCAM/1159]


| Catalog No        | Format                 | Size   |
|-------------------|------------------------|--------|
|                   | Bioreactor Concentrate | 0.1 ml |
| RT171577-MSM2-BC1 | Bioreactor Concentrate | 0.5 ml |

| Applications               | Tested Dillution | Note  |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml         | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |


| Product Details        |  |  |
|------------------------|--|--|
| Clone                  | EpCAM/1159   |  |
| Gene Name              | EPCAM  |  |
| Immunogen              | Recombinant full-length Rat Epcam protein            |  |
| Host                   | Mouse  |  |
| Clonality              | Monoclonal   |  |
| Isotype / Light Chain  | IgG1 / Kappa   |  |
| Mol. Weight of Antigen | 40-43kDa   |  |
| Cellular Localization  | Cell junction, Lateral cell membrane, Tight junction |  |
| Species Reactivity     | Mouse, Rat   |  |
| Positive Control       | Recombinant full-length rat Epcam protein.           |  |

<sup>\*</sup>Optimal dilution for a specific application should be determined.

#### Product Images for Ep-CAM / CD326 (Rat) Antibody



Formalin-fixed, paraffin-embedded Rat Lung stained with Ep-CAM Monoclonal Antibody (Epcam/1159).



Formalin-fixed, paraffin-embedded Rat Colon stained with Ep-CAM Monoclonal Antibody (Epcam/1159).

### **Specificity & Comments**

Epitopes of Epcam/1158 and Epcam/1159 monoclonal antibodies are different. EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas.

### **Supplied As**

Bioreactor Concentrate with 0.05% Azide. Contact us if you require it in a different format.

#### Storage and Stability

Store at 2 to 8°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

## **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

