

CD99 / MIC2 (Ewing's Sarcoma Marker) Antibody

Catalog No	Format	Size
4267-MSM4-CS0	Culture Supernatant	0.1 ml
4267-MSM4-CS1	Culture Supernatant	0.5 ml


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Gene Name	CD99
Immunogen	Human acute lymphocytic leukemia T-cells (12E7); Recombinant human MIC2 protein (MIC2/877)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	27-32kDa
Cellular Localization	Cell surface
Species Reactivity	Human
Positive Control	MOLT-4 cells. Pancreas or Ewing's sarcoma.

**Optimal dilution for a specific application should be determined.*

Product Images for CD99 / MIC2 (Ewing's Sarcoma Marker) Antibody



Specificity & Comments

Supplied As

Tissue culture supernatant with 0.05% Azide. Contact us if you require it in a different format.

Storage and Stability

Store at 2 to 8°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Immunology

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.