

## CD63 (Late Endosomes Marker) Antibody

Mouse Monoclonal Antibody [Clone MX-49.129.5]

Catalog No	Format	Size
------------	--------	------


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	

### Product Details


<b>Clone</b>	MX-49.129.5
<b>Gene Name</b>	CD63
<b>Immunogen</b>	Full length CD63 of human origin
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	26kDa (core protein); 30-60kDa (glycosylated)
<b>Cellular Localization</b>	Cytoplasm
<b>Species Reactivity</b>	Human, Mouse
<b>Positive Control</b>	HL60 SK-MEL-28 THP-1 or NIH/3T3 cells. Melanoma or lymphoma.

\*Optimal dilution for a specific application should be determined.

### Product Images for CD63 (Late Endosomes Marker) Antibody



Flow Cytometric staining of CD63 on human PBMC cells. Black: cells alone; Green: Isotype Control; Red: PE-labeled CD63 Monoclonal Antibody (MX-49.129.5).



Flow Cytometric staining of Mouse CD63 on NIH/3T3 cells. Black: cells alone; Green: Isotype Control; Red: PE-labeled CD63 Monoclonal Antibody (MX-49.129.5).

### **Specificity & Comments**

This MAAb recognizes protein of 26kDa-60kDa, which is identified as CD63. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.

---

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### **Supplied As**

Antibody Purified from Bioreactor Concentrate by Protein A/G and conjugated to various reporter molecules. Prepared in 10mM PBS with 0.05% BSA and 0.05% azide. Contact us if you require this Ab in a different format.

---

### **Storage and Stability**

Antibody with azide - store at 4 to 8°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

### **Research Areas**

Immunology

---