

Recombinant Double Stranded DNA (dsDNA) (Nuclear Marker) Antibody

Rabbit Monoclonal Antibody [Clone DSD/8204R]

Catalog No	Format	Size
RBM7-8204-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
RBM7-8204-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
RBM7-8204-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	DSD/8204R
Gene Name	
Immunogen	Nuclei of Burkitt's cells
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	Not Known
Cellular Localization	Nucleus.
Species Reactivity	Human
Positive Control	Jurkat or HeLa cells. Human tonsil or colon. Raji

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Double Stranded DNA (dsDNA) (Nuclear Marker) Antibody



Formalin-fixed, paraffin-embedded human placenta stained with dsDNA Recombinant Rabbit Monoclonal Antibody (DSD/8204R). HIER: Tris/EDTA, pH9.0, 45min. 2: HRP-polymer, 30min. DAB, 5min.



Formalin-fixed, paraffin-embedded human ovarian cancer (serous) stained with dsDNA Recombinant Rabbit Monoclonal Antibody (DSD/8204R). HIER: Tris/EDTA, pH9.0, 45min. 2: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

This monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This MAb recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This MAb produces a homogeneous staining pattern in the nucleus of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3',5'-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
