

# **Recombinant c-Myc Oncoprotein Antibody**

Rabbit Monoclonal Antibody [Clone MYC/7682R]

Catalog No	Format	Size
4609-RBM12-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4609-RBM12-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4609-RBM12-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

MYC/7682R	
MYC	
Recombinant fragment (around aa1-200) of human MYC protein (exact sequence is proprietary)	
Rabbit	
Monoclonal	
IgG / Kappa	
62-64kDa	
Nucleus.	
Human	
HL-60 cells. Human cervical carcinoma or heart tissue.	

<sup>\*</sup>Optimal dilution for a specific application should be determined.

### **Product Images for Recombinant c-Myc Oncoprotein Antibody**

#### **Specificity & Comments**

It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.Â?

#### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

# Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

# **Research Areas**

Apoptosis, Autophagy, Breast Cancer, Cardiovascular, Developmental Biology, Immunology, Bladder Cancer, Colon Cancer, Cytokine Signaling, Infectious Disease, Lung Cancer, MAPK Signaling, Nuclear Marker, Signal Transduction, Stem Cell Differentiation, Transcription Factors

## **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

