

# Recombinant PDCD1 / PD1 / CD279 (Programmed Cell Death 1) Antibody

Rabbit Monoclonal Antibody [Clone PDCD1/8170R]

Catalog No	Format		Size
5133-RBM17-P0	Purified Ab with BSA and Azi	ide at 200ug/ml	20 ug
5133-RBM17-P1	Purified Ab with BSA and Azi	ide at 200ug/ml	100 ug
5133-RBM17-P1ABX	Purified Ab WITHOUT BSA a	and Azide at 1.0mg/ml	100 ug
Applications	Tested Dillution	Note	

Applications	Testeu Dillution	Note
Immunohistochemistry (IHC)	5	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C
		followed by cooling at RT for 20 minutes

Product Details		
Clone	PDCD1/8170R	
Gene Name	PDCD1	
Immunogen	Recombinant fragment (around aa1-200) of human PDCD1 protein (exact sequence is proprietary)	
Host	Rabbit	
Clonality	Monoclonal	
Isotype / Light Chain	IgG / Kappa	
Mol. Weight of Antigen	55kDa	
Cellular Localization	Cell Surface. Cytoplasm.	
Species Reactivity	Human	
Positive Control	Human tonsil lymph node or appendix.	

\*Optimal dilution for a specific application should be determined.

Product Images for Recombinant PDCD1 / PD1 / CD279 (Programmed Cell Death 1) Antibody

### **Specificity & Comments**

PDCD-1 (programmed cell death-1 protein), also designated CD279, is a type I transmembrane receptor and a member of the immunoglobin gene superfamily. It is expressed on activated T-cells, B-cells, and myeloid cells. Anti-PDCD-1 is a marker of angioimmunoblastic lymphoma and suggests a unique cell of origin for this neoplasm. Unlike CD10 and BCL6, PDCD-1 is expressed by few B-cells, so anti-PDCD-1 may be a more specific and useful diagnostic marker in angioimmunoblastic lymphoma. In addition, PDCD-1 expression provides evidence that angioimmunoblastic lymphoma is a neoplasm derived from germinal center-associated T-cells.

## Supplied As

200ug/ml of Ab purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

#### Storage and Stability

Antibody with azide - store at 2 to  $8^{\circ}$ C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

#### **Research Areas**

Cardiovascular, Immunology, B Cell Markers, Infectious Disease, PD-1 blockade immunotherapy

#### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

