

Recombinant CD45 / LCA (B-Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rPTPRC/1147]

Catalog No	Format	Size
5788-MSM46-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5788-MSM46-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5788-MSM46-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	rPTPRC/1147
Gene Name	PTPRC
Immunogen	Recombinant full-length human PTPRC protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	180-220kDa
Cellular Localization	Cell Surface. Cytoplasm.
Species Reactivity	Human
Positive Control	Jurkat Ramos U-698 or GA-10 cells. Human tonsil.

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant CD45 / LCA (B-Cell Marker) Antibody



Western blot analysis of human spleen tissue lysate using CD45 Recombinant Mouse Monoclonal Antibody (rPTPRC/1147).



Formalin-fixed, paraffin-embedded human spleen stained with CD45 Recombinant Mouse Monoclonal Antibody (rPTPRC/1147). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology, Immunology, B Cell Markers, Dendritic Cell Marker, Hematopoietic Stem Cells, Mesenchymal Stem Cell Differentiation
