

BATF2 / SARI (Transcription Factor) Antibody

Mouse Monoclonal Antibody [Clone PCR-P-BATF2-2B9]

Catalog No	Format	Size
116071-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
116071-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
116071-MSM2-P1ABX	Purified Ab WITHOUT BSA at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	PCR-P-BATF2-2B9
Gene Name	BATF2
Immunogen	Recombinant fragment (around aa5-140) of human BATF2 (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	29.4kDa
Cellular Localization	Nucleus.
Species Reactivity	Human
Positive Control	colon or pancreas. Human spleen

*Optimal dilution for a specific application should be determined.

Product Images for BATF2 / SARI (Transcription Factor) Antibody



Flow cytometric analysis of PFA-fixed HeLa cells. BATF2 Mouse Monoclonal Antibody (PCR-P-BATF2-2B9) followed by goat anti-mouse IgG-CF488 (blue); isotype control (red).



Formalin-fixed, paraffin-embedded human prostate carcinoma stained with BATF2 Mouse Monoclonal Antibody (PCR-P-BATF2-2B9). Inset: PBS instead of primary antibody; secondary only negative control.




Analysis of Protein Array containing more than 19,000 full-length human proteins using BATF2 Mouse Monoclonal Antibody (PCR-BATF2-2B9). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Specificity & Comments

Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II cis-regulatory region sequence-specific DNA binding activity. Predicted to be involved in defense response to protozoan; myeloid dendritic cell differentiation; and regulation of transcription by RNA polymerase II. Predicted to be part of chromatin. Predicted to be active in nucleus. This antibody recognizes a transcription factor involved in differentiation of CD8+ thymic dendritic cells. BATF2 has been implicated in breast cancer, malignant glioma and metastasis melanoma progression.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.



SDS-PAGE Analysis of Purified BATF2 Mouse Monoclonal Antibody (PCR-BATF2-2B9). Confirmation of Purity and Integrity of Antibody.

Supplied As

200µg/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.