

## Recombinant CD162 (Selectin P Ligand) Antibody

Rabbit Monoclonal Antibody [Clone PSGL1/8192R]

| Catalog No      | Format  | Size   |
|-----------------|---|--------|
| 6404-RBM5-P0    | Purified Ab with BSA and Azide at 200ug/ml    | 20 ug  |
| 6404-RBM5-P1    | Purified Ab with BSA and Azide at 200ug/ml    | 100 ug |
| 6404-RBM5-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications               | Tested Dillution | Note  |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml         | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

### Product Details


|                        |  |
|------------------------|--|
| Clone                  | PSGL1/8192R  |
| Gene Name              | SELPLG   |
| Immunogen              | Recombinant full-length human SELPLG (CD162) protein |
| Host                   | Rabbit   |
| Clonality              | Monoclonal   |
| Isotype / Light Chain  | IgG / Kappa  |
| Mol. Weight of Antigen | 43-120kDa  |
| Cellular Localization  | Cell surface.  |
| Species Reactivity     | Human  |
| Positive Control       | Human tonsil or spleen.                              |

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant CD162 (Selectin P Ligand) Antibody




Formalin-fixed, paraffin-embedded human spleen stained with CD162 Recombinant Rabbit Monoclonal Antibody (PSGL1/8192R). HIER: Tris/EDTA, pH9.0, 45min. 2: HRP-polymer, 30min. DAB, 5min.



IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using PSGL1/8192R at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified CD162 Recombinant Rabbit Monoclonal Antibody (PSGL1/8192R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human tonsil stained with CD162 Recombinant Rabbit Monoclonal Antibody (PSGL1/8192R). HIER: Tris/EDTA, pH9.0, 45min. 2: HRP-polymer, 30min. DAB, 5min.

### Specificity & Comments

CD162 glycoprotein functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L- selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking during inflammation by tethering of leukocytes to activated platelets or endothelia expressing selectins. This protein requires two post-translational modifications, tyrosine sulfation and the addition of the sialyl Lewis x tetrasaccharide (sLex) to its O-linked glycans, for its high-affinity binding activity. Aberrant expression of this gene and polymorphisms in this gene are associated with defects in the innate and adaptive immune response.Â ?

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Complement System