

Recombinant Human Nuclear Antigen (HNA) (Human Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone 235-1R]


Catalog No	Format	Size

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details		
Clone	235-1R	
Gene Name		
Immunogen	Nuclei of human myeloid leukemia biopsy cells	
Host	Rabbit	
Clonality	Monoclonal	
Isotype / Light Chain	IgG / Kappa	
Mol. Weight of Antigen	70kDa & 80kDa	
Cellular Localization	Nucleus.	
Species Reactivity	Human	
Positive Control	All human cells. Human tonsil.	

^{*}Optimal dilution for a specific application should be determined.

Product Images for Recombinant Human Nuclear Antigen (HNA) (Human Cell Marker) Antibody



SDS-PAGE Analysis of Purified HuNu Recombinant Rabbit Monoclonal Antibody (235-1R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. It is a part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. MAb 235-1R recognizes an antigen associated with the nuclei in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in subcellular fractions. It produces a speckled pattern in normal and malignant cells and may be used to stain the nuclei of cells in fixed or frozen tissue sections. It can also be used with paraformaldehyde fixed frozen tissue or cell preparations.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

