

ZNF232 (Transcription Factor) Antibody

Mouse Monoclonal Antibody [Clone PCR-P-ZNF232-2B3]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 7775-MSM3-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 7775-MSM3-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 7775-MSM3-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|---------------------|---|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

Product Details


| | |
|-------------------------------|--|
| Clone | PCR-P-ZNF232-2B3 |
| Gene Name | ZNF232 |
| Immunogen | Recombinant fragment (around aa29-109) of human ZNF232 protein |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG2a |
| Mol. Weight of Antigen | 47.69kDa |
| Cellular Localization | Nucleus. |
| Species Reactivity | Human |
| Positive Control | HeLa or U87 cells. |

*Optimal dilution for a specific application should be determined.


Product Images for ZNF232 (Transcription Factor) Antibody



Flow cytometric analysis of PFA-fixed HeLa cells. ZNF232 Mouse Monoclonal Antibody (PCR-P-ZNF232-2B3) followed by goat anti-mouse IgG-CF488 (blue), unstained cells (red).



Analysis of Protein Array containing more than 19,000 full-length human proteins using ZNF232 Mouse Monoclonal Antibody (PCR-P-ZNF232-2B3). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human tumor of unknown origin stained with ZNF232 Mouse Monoclonal Antibody (PCRP-ZNF232-2B3). HIER: Tris/EDTA, pH9.0, 45min. 2°C; HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II cis-regulatory region sequence-specific DNA binding activity. Predicted to be involved in regulation of transcription by RNA polymerase II. Located in cytosol and nucleoplasm. [provided by Alliance of Genome Resources, Apr 2022]May be involved in transcriptional regulation.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
