

## FGF23 (Fibroblast Growth Factor 23) Antibody

Mouse Monoclonal Antibody [Clone FGF23/6419]

Catalog No	Format	Size
8074-MSM29-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
8074-MSM29-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
8074-MSM29-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details


<b>Clone</b>	FGF23/6419
<b>Gene Name</b>	FGF23
<b>Immunogen</b>	Recombinant fragment (around aa1-251) of human FGF23 protein (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	12-32kDa
<b>Cellular Localization</b>	Secreted (extracellular)
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Human kidney tissue.

\*Optimal dilution for a specific application should be determined.

### Product Images for FGF23 (Fibroblast Growth Factor 23) Antibody



SDS-PAGE Analysis of Purified FGF23 Mouse Monoclonal Antibody (FGF23/6419). Confirmation of Purity and Integrity of Antibody.



IHC analysis of formalin-fixed, paraffin-embedded human kidney stained using FGF23/6419 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

### Specificity & Comments

Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast growth factor-2 (FGF-2), also designated basic FGF, are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10 through FGF-23. Members of the FGF family share 30-55% amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family, including four tyrosine kinases designated Flg (FCFR 1), Bok (FCFR L), TKF and FCFR 3.

2 Union Square, Union City, CA 94587, Tel: 1-866-2-NEOBIO (1-866-263-6246)  
orders@NeoBiotechnologies.com, www.NeoBiotechnologies.com

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

AKT Signaling, Breast Cancer, Cardiovascular, Infectious Disease, MAPK Signaling, Signal Transduction

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---