

# Recombinant Biotin (Vitamin B7 or Vitamin H) Antibody

Mouse Monoclonal Antibody [Clone rBTN/8819]


Catalog No	Format	Size
MSM8-8819-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM8-8819-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM8-8819-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	


Product Details		
Clone	rBTN/8819	
Gene Name		
Immunogen	Biotinylated sheep immunoglobulin	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG1 / Kappa	
Mol. Weight of Antigen	244Da	
Cellular Localization	Depends on the target	
Species Reactivity	All species	
Positive Control	Biotinylated proteins in solution or on tissues.	

<sup>\*</sup>Optimal dilution for a specific application should be determined.

## Product Images for Recombinant Biotin (Vitamin B7 or Vitamin H) Antibody







Formalin-fixed, paraffin-embedded human kidney stained with Biotinylated Lambda Light Chain probe followed by Anti-Biotin Recombinant Mouse Monoclonal Antibody (rBTN/8819).

### **Specificity & Comments**

It recognizes both the free and protein-conjugated (either soluble or cell bound) form of biotin. This MAb is highly specific to biotin and shows no cross-reaction with other structurally related compounds. It has a very high affinity for biotin and is excellent for use in various amplification techniques. In some applications, localization of biotinylated probes with avidin produces unacceptably high background staining. Anti-biotin antibody may be substituted to

#### **Supplied As**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

#### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-





## **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

