

## Recombinant IL3RA / CD123 (Acute Myeloid Leukemia Marker) Antibody

Mouse Monoclonal Antibody [Clone rIL3RA/9212]

Catalog No	Format	Size
3563-MSM8-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3563-MSM8-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3563-MSM8-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details


<b>Clone</b>	rIL3RA/9212
<b>Gene Name</b>	IL3RA
<b>Immunogen</b>	Recombinant fragment of human IL3RA protein (around aa 26-171) (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2b / Kappa
<b>Mol. Weight of Antigen</b>	~70kDa
<b>Cellular Localization</b>	Cell Surface. Cytoplasm.
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	lymph node or stomach RPMI-8226 or HDLM-2 cells. Human tonsil THP-1

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant IL3RA / CD123 (Acute Myeloid Leukemia Marker) Antibody



SDS-PAGE Analysis of Purified IL3RA / CD123 Recombinant Mouse Monoclonal Antibody (rIL3RA/9212). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human tonsil stained with IL3RA / CD123 Recombinant Mouse Monoclonal Antibody (rIL3RA/9212). Inset: PBS instead of primary antibody; secondary only negative control.

### Specificity & Comments

CD123 is an interleukin 3 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL3 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL3. This gene and the gene encoding the colony stimulating factor 2 receptor alpha chain (CSF2RA) form a cytokine receptor gene cluster in a X-Y pseudo-autosomal region on chromosomes X or Y.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Immunology, AKT Signaling, Cytokine Signaling, Hematopoietic Stem Cells, Signal Transduction

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---