

CD9 (TSPAN29) (Motility-Related Protein-1) Antibody

Mouse Monoclonal Antibody [Clone CD9/7416]

Catalog No	Format	Size
928-MSM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
928-MSM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
928-MSM5-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	CD9/7416
Gene Name	CD9
Immunogen	Recombinant full-length human CD9 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	24kDa
Cellular Localization	Cell Surface. Cytoplasm.
Species Reactivity	Human
Positive Control	BT-20 HeLa or ZR-75 cells. Human cervix or esophagus, Brain

*Optimal dilution for a specific application should be determined.


Product Images for CD9 (TSPAN29) (Motility-Related Protein-1) Antibody




Formalin-fixed, paraffin-embedded human lymph node stained with CD9 Mouse Monoclonal Antibody (CD9/7416). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified CD9 Mouse Monoclonal Antibody (CD9/7416). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Brain tissue lysate using CD9 Mouse Monoclonal Antibody (CD9/7416).



Formalin-fixed, paraffin-embedded human pancreas stained with CD9 Mouse Monoclonal Antibody (CD9/7416). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

CD9 is a type IV transmembrane glycoprotein with four transmembrane domains. CD9 on pre-B cells may play a role in cell-cell adhesion. In addition, CD9 may play a role in signal transduction mediated by interaction with low molecular weight GTP binding proteins. CD9 is expressed on early B cells, eosinophils, basophils and activated T cells and is a major component of the platelet cell surface. It is also expressed on most non-T acute lymphoblastic leukemia cells and on some acute myeloid and chronic lymphoid leukemia.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Endothelial Cell Marker, Infectious Disease

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.