

Parvalbumin / PVALB Antibody

Mouse Monoclonal Antibody [Clone PVALB/7601]

Catalog No	Format	Size
5816-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5816-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5816-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	PVALB/7601
Gene Name	PVALB
Immunogen	Recombinant fragment (around aa1-110) of human PVALB protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	12kDa
Cellular Localization	Nucleus. Cell junctions.
Species Reactivity	Human
Positive Control	Human brain or kidney.

*Optimal dilution for a specific application should be determined.

Product Images for Parvalbumin / PVALB Antibody



Formalin-fixed, paraffin-embedded human kidney stained with PVALB Mouse Monoclonal Antibody (PVALB/7601). Inset: PBS instead of primary antibody; secondary only negative control.



SDS-PAGE Analysis of Purified PVALB Mouse Monoclonal Antibody (PVALB/7601). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

The family of EF-hand type Ca²⁺-binding proteins includes calbindin (previously designated vitamin D-dependent Ca²⁺-binding protein), S-100 β and γ , calgranulins A (also designated MRP8), B (also designated MRP14) and C (S-100 like proteins) and the parvalbumin family members, including parvalbumin β and parvalbumin γ , also designated oncomodulin (OCM). Structurally and evolutionarily conserved, parvalbumin β and OCM proteins are distinct in expression and function. Parvalbumin β , also designated parvalbumin (PV), is most abundantly expressed in fast-contracting muscles with lower expression levels in brain and some endocrine tissues, including kidney and parathyroid. Research indicates that parvalbumin β plays a significant role in muscle relaxation. OCM was originally thought to have expression restricted to neoplastic tissues, early embryonic cells and certain tumor cell lines. Recent research shows that OCM is also expressed and secreted by macrophages where, in the retina it binds to retinal ganglion cells (RGCs) and functions to promote axon regeneration. OCM has also been detected in the auditory sensory cells of the organ of Corti in mammals. In humans, two different loci on chromosome 7 have been identified as OCM and OCM-like (LOC4951). These genes encode proteins 109 amino acids in length which share 99% sequence identity.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Nuclear Marker, Transcription Factors
