

Recombinant Ki-67 (Proliferating Cell Marker) Antibody


Mouse Monoclonal Antibody [Clone rMKI67/9616]

Catalog No	Format	Size
4288-MSM35-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4288-MSM35-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4288-MSM35-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note	
Product Details			
Clone	rMKI67/9616		
Gene Name	MKI67		
Immunogen	Recombinant fragment (around aa2200-2500) of human MKI67 protein (exact sequence is proprietary)		
Host	Mouse		
Clonality	Monoclonal		
Isotype / Light Chain	IgG2b / Kappa		
Mol. Weight of Antigen	345kDa and 395kDa		
Cellular Localization	Nucleus.		
Species Reactivity	Human	Human	
Positive Control	Any actively proliferating cells. Human	Any actively proliferating cells. Human skin, tonsil or lymph node.	

^{*}Optimal dilution for a specific application should be determined.

Product Images for Recombinant Ki-67 (Proliferating Cell Marker) Antibody



SDS-PAGE Analysis of Purified Ki-67 Recombinant Mouse Monoclonal Antibody (rMKl67/9616). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Ki-67 antigen is a nuclear, non-histone protein that is present in all stages of the cell cycle except G0. This characteristic makes Ki-67 an excellent marker for proliferating cells and is commonly used as one of the prognostic factors in cancer studies. A correlation has been demonstrated between Ki-67 index and the histo-pathological grade of neoplasms. Assessment of Ki-67 expression in renal and ureter tumors shows a correlation between tumor proliferation and disease progression, thus making it possible to differentiate high-risk patients. Ki-67 expression may also prove to be important for distinguishing between malignant and benign peripheral nerve sheath tumors. Ki-67 labeling index has been shown to be a prognostic marker in a number of neoplasms including grade II astrocytoma, oligodendroglioma, colon carcinoma, and breast carcinoma. In general, Ki-67 is a good marker of proliferating cell populations. ?

Supplied As

200ug/ml of Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Neuroscience, Nuclear Marker

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product