

Recombinant Myelin Basic Protein Antibody

Mouse Monoclonal Antibody [Clone rMBP/9431]


Catalog No	Format	Size
4155-MSM13-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4155-MSM13-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4155-MSM13-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details		
Clone	rMBP/9431	
Gene Name	MBP	
Immunogen	Recombinant fragment (around aa1-100) of human MBP protein (exact sequence is proprietary)	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG2b / Kappa	
Mol. Weight of Antigen	13-21kDa	
Cellular Localization	Cell membrane, Myelin membrane	
Species Reactivity	Human	
Positive Control	Human brain or astrocytoma tissue.	

^{*}Optimal dilution for a specific application should be determined.

Product Images for Recombinant Myelin Basic Protein Antibody





Formalin-fixed, paraffin-embedded human brain stained with Myelin Basic Protein Recombinant Mouse Monoclonal Antibody (rMBP/9431). Inset: PBS instead of primary antibody; secondary only negative control.

Western Blot Analysis of human brain tissue lysate using Myelin Basic Protein Recombinant Mouse Monoclonal Antibody (rMBP/9431).

Specificity & Comments

Myelin basic protein (MBP) is the second most abundant protein in central nervous system (CNS) myelin: it comprises 30% of the total protein and about 10% of the dry weight of myelin. It is the only structural protein found so far to be essential for formation of CNS myelin, and has been called the executive molecule of myelin . MBP can interact with a number of polyanionic proteins including actin, tubulin, calmodulin, and clathrin, and negatively charged lipids, and acquires structure on binding to them. It may act as a membrane actin-binding protein, which might allow it to participate in transmission of extracellular signals to the cytoskeleton in oligodendrocytes and tight junctions in myelin. MBP may be applicable as a marker for oligodendrogliomas. rMBP/9301 is used in clinical diagnostics to detect MBP levels or myelination in human MBP. rMBP/9301 is used in a two-site ELISA with clone MBP/4273. Both are used interchangeably as capture and detection antibody.

Supplied As

200ug/ml of Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Complement System, Developmental Biology, Neuroscience

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

